

ASIA-ICHC Selected Quality Measures  
 UDS Health Center Comparison Report 2017  
 last updated 8/2018  
 (shared at <http://www.asiaohio.org>)

ASIA – ICHC strives to serve the community with high quality of patient care. Combined across all sites.

The following data shows ICHC across all sites compared with the national average. Regional information can be found via the following link for *Better Health Greater Cleveland* data:

<http://www.betterhealthpartnership.org>.

<b>Target Measure</b>	<b>Description</b>	<b>UDS 2017 ICHC %</b>	<b>National Average</b>
Diabetes A1C	Percentage of diabetic patients whose HbA1c levels are greater than 9 percent.	27.61%	32.95%
Cardiovascular	Percentage of adult patients with diagnosed hypertension whose most recent blood pressure was less than 140/90.	67.47%	62.71%
Cancer – Female	Percentage of women 21-64 years of age who received one or more tests to screen for cervical cancer during the measurement year or two years prior to the measurement year.	45.44%	55.67%
Child Health	Percentage of children who have received age appropriate vaccines on or before their 2nd birthday.	0.00%	40.24%
Behavioral Health	Percentage of patients aged 12 years and older screened for clinical depression on the date of the visit using an age-appropriate standardized depression screening tool, and, if screening is positive, for whom a follow-up plan is documented on the date of the positive screen.	57.66%	66.15%
Weight Assessment and Counseling for Children and Adolescents	Percentage of patients age 2 to 17 years who had a visit during the current year and who had Body Mass Index (BMI) Percentile documentation, counseling for nutrition, and counseling for physical activity during the measurement year.	40.71%	65.85%
Adult Weight Screening and Follow-Up	Percentage of patients age 18 years or older who had their <i>Body Mass Index (BMI) calculated at the last visit or within the last six months and, if they were overweight or underweight, had a follow-up plan documented.</i>	44.76%	63.85%
Tobacco Use Screening and Cessation (Tobacco Use Assessment and Cessation Counseling)	Percentage of patients age 18 years and older who were screened for tobacco use at least once during the measurement year or prior year AND who received cessation counseling intervention and/or pharmacotherapy if identified as a tobacco user..	87.17%	87.50%

ASIA-ICHC Selected Quality Measures  
 UDS Health Center Comparison Report 2017  
 last updated 8/2018  
 (shared at <http://www.asiaohio.org>)

Coronary Artery Disease (CAD): Lipid Therapy	Percentage of patients age 18 years and older with a <i>diagnosis of CAD prescribed a lipid lowering therapy</i> (based on current ACC/AHA guidelines) during the measurement year.	53.85%	70.72%
Ischemic Vascular Disease (IVD): Aspirin Therapy	Percentage of patients age 18 years and older who were discharged alive for <i>acute myocardial infarction (AMI), coronary artery bypass graft (CABG), or percutaneous transluminal coronary angioplasty (PTCA), or who had a diagnosis of Ischemic Vascular Disease (IVD), and who had documentation of use of aspirin or another antithrombotic</i> during the measurement year.	55.56%	79.27%
Colorectal Cancer Screening	Percentage of patients age 50 to 75 years who had appropriate <i>screening for colorectal cancer</i> (includes colonoscopy ≤ 10 years, flexible sigmoidoscopy ≤ 5 years, or annual fecal occult blood test)	35.80%	42.02%
Referral Tracking	Percentage of specialist referrals that are completed and consult notes reviewed by providers.	75.27% (Referral Data for Jan 2018)	-
Patient Satisfaction Survey	Percentage of patient evaluation of the experience as very satisfied.	69.23%	-